Distribution: Viet Nam.

Type: Holotype, female, Dalat, Dralac Prov., Viet Nam, 12–14 November 1959, N. R. Spencer, light trap (Type no. 71178, USNM).

Discussion: Pseudostilobezzia appears to be closely related to Stilobezzia, subgenus Stilobezzia, as evident from the general appearance, wing venation, head and antennal structure, but differs in the condition of the tarsi and claws. In females of Stilobezzia the fifth tarsomeres bear strong ventral spines if the wing lacks macrotrichia (subgenus Stilobezzia), or if spines are lacking the wing bears macrotrichia (subgenus Neostilobezzia). In Stilobezzia the female tarsal claws are large and unequal on all legs, or reduced to a single long claw with a small basal barb. In the genus Monohelea, which differs considerably in general features and wing venation from Pseudostilobezzia, the female tarsal claws are sometimes small and equal on the fore and mid legs, but are very unequal, or reduced to a single long claw with a basal barb, on the hind leg.

We are very pleased to name this species in honor of Dr. H. Elliott McClure of the Migratory Animal Pathological Survey, Bangkok, Thailand, in appreciation of his long and enthusiastic help and guidance in our studies of Southeast Asian Ceratopogonidae.

REFERENCE

Das Gupta, S. K., and W. W. Wirth. 1968. Revision of the Oriental Species of Stilobezzia Kieffer (Diptera, Ceratopogonidae). Bull. U. S. Nat. Mus. 283: 1–149.


STUDIES ON IDIOCERINAE LEAFHOPPERS: X. IDIOSCOPUS NITIDULUS (WALKER), NEW COMBINATION (HOMOPTERA: CICADELLIDAE)

J. Maldonado-Capriles

Department of Biology, University of Puerto Rico, Mayaguez, Puerto Rico 00708

ABSTRACT—*Idiocerus nitidulus* Walker is transferred to the genus *Idioscopus* Baker, and the male genitalia are illustrated.

Among specimens recently obtained for study from the Leiden Museum, The Netherlands, there were some that I suspected to be *Idiocerus nitidulus* Walker 1870 because they were collected in Java. Dr. W. J. Knight at the British Museum (N.H.) kindly agreed to compare a female and a male with Walker's type. After examining



Idioscopus nitidulus (Walker), male. 1, genital capsule, lateral view; 2, acdeagus, caudal view; 3, same, lateral view; 4, style, ventral view; 5, same, lateral view.

the specimens he reported, "I have compared your female specimens of *Idiocerus nitidulus* Walker with the type and although it is a female and the type is a male they are identical externally". "The genitalia of your male are identical although much smaller than the type which

is approximately the same size as your female specimen".

In my paper on the Indian and Philippine species of Idiocerus (Maldonado-Capriles 1964) I moved to Idioscopus Baker several species of the latter. On page 93 I wrote, about Idioscopus incertus (Baker), "Very close to Idioscopus niveosparsus, perhaps only a subspecies or an extreme variety". Idioscopus niveosparsus Lethierry 1889 is very variable in coloration, judging from the many specimens studied before and the few at hand; this is to be expected of a very prolific species with such a wide range of distribution. The genitalia of I. nitidulus are very close to those of I. niveosparsus and I. incertus. The seventh sternum of the female is similar to that of niveosparsus. Thus, probably I. nitidulus is also a subspecies or just another extreme form of niveosparsus. Unfortunately, nitidulus has priority over niveosparsus. Because I. niveosparsus is of economic importance the name has been used in many papers and, therefore, a change of name would cause much inconvenience. Therefore, at the present time I am only moving nitidulus to the correct genus and leaving the problem of priority until more evidence, preferably biological, can be obtained to clearly establish the synonymy. In such event the name niveosparsus should be conserved.

Idioscopus nitidulus (Walker), new combination

Idiocerus nitidulus Walker 1870. Linn. Soc. London Jour. Zool. 10:322

Study of the male type of this species shows that it properly belongs in *Idioscopus*. The genitalia of a specimen labeled J. Sonneveldt, Makasser, compared with that of the type, are illustrated in figures 1 to 5. A female labeled Tijomas, W. Java, has been also compared with the type. Both specimens are in the Leiden Museum. This species runs to *I. incertus* and *I. niveosparsus* in my paper of 1964.

REFERENCE

Maldonado-Capriles, J. 1964. Studies on Idiocerinae leafhoppers: II. The Indian and Philippine species of *Idiocerus* and the genus *Idioscopus* (Homoptera: Cicadellidae). Proc. Entomol. Soc. Wash. 66;89–100.

TWO NEW SPECIES OF ANTS OF THE GENERA TAPINOMA FOERSTER AND PARATRECHINA MOTSCHOULSKY FROM PUERTO RICO (HYMENOPTERA: FORMICIDAE)

DAVID R. SMITH

Systematic Entomology Laboratory, Agricultural Research Service, USDA, c/o U. S. National Museum, Washington, D. C. 20560

ROBERT J. LAVIGNE

Entomology Section, University of Wyoming, Laramie, Wyoming 82070

ABSTRACT—Two new species of ants from Puerto Rico are described, the worker and male of *Tapinoma* resenum, n. sp., and each caste of Paratrechina (*Nylanderia*) cisipa, n. sp.

No new species of ants have been described from Puerto Rico since M. R. Smith (1936) described *Prenolepis microps* (a species now placed in the genus *Paratrechina*). He listed and keyed 66 species of ants and referred to the previous literature concerning ants of Puerto Rico. Wolcott (1948) summarized the available information on their habits, and provided additional locality records.

During a year's sojourn (1969–1970) in Puerto Rico while working for the Puerto Rico Nuclear Center, the junior author had the opportunity to collect ants, along with other insects, at various localities around the island. Among the ants collected were 2 species not only new to Puerto Rico, but so far as we can determine, undescribed. These species, representing a new *Paratrechina* and a new *Tapinoma*, are described herein.